



PAIN

Volume 152, Issue 3, Supplement, March 2011, Pages S2-S15

Review

Central sensitization: Implications for the diagnosis and treatment of pain

Clifford J. Woolf

Show more

<https://doi.org/10.1016/j.pain.2010.09.030>

[Get rights and content](#)

Abstract

Nociceptor inputs can trigger a prolonged but reversible increase in the excitability and synaptic efficacy of neurons in central nociceptive pathways, the phenomenon of central sensitization. Central sensitization manifests as pain hypersensitivity, particularly dynamic tactile allodynia, secondary punctate or pressure hyperalgesia, aftersensations, and enhanced temporal summation. It can be readily and rapidly elicited in human volunteers by diverse experimental noxious conditioning stimuli to skin, muscles or viscera, and in addition to producing pain hypersensitivity, results in secondary changes in brain activity that can be detected by electrophysiological or imaging techniques. Studies in clinical cohorts reveal changes in pain sensitivity that have been interpreted as revealing an important contribution of central sensitization to the pain phenotype in patients with fibromyalgia, osteoarthritis, musculoskeletal disorders with generalized pain hypersensitivity, headache, temporomandibular joint disorders, dental pain, neuropathic

hypersensitivity, headache, temporomandibular joint disorders, dental pain, neuropathic pain, visceral pain hypersensitivity disorders and post-surgical pain. The comorbidity of those pain hypersensitivity syndromes that present in the absence of inflammation or a neural lesion, their similar pattern of clinical presentation and response to centrally acting analgesics, may reflect a commonality of central sensitization to their pathophysiology. An important question that still needs to be determined is whether there are individuals with a higher inherited propensity for developing central sensitization than others, and if so, whether this conveys an increased risk in both developing conditions with pain hypersensitivity, and their chronification. Diagnostic criteria to establish the presence of central sensitization in patients will greatly assist the phenotyping of patients for choosing treatments that produce analgesia by normalizing hyperexcitable central neural activity. We have certainly come a long way since the first discovery of activity-dependent synaptic plasticity in the spinal cord and the revelation that it occurs and produces pain hypersensitivity in patients. Nevertheless, discovering the genetic and environmental contributors to and objective biomarkers of central sensitization will be highly beneficial, as will additional treatment options to prevent or reduce this prevalent and promiscuous form of pain plasticity.



[Previous article](#)

[Next article](#)



Choose an option to locate/access this article:

Check if you have access through your login credentials or your institution.

[Check Access](#)

[Recommended articles](#)

[Citing articles \(0\)](#)

Central sensitization: implications for the diagnosis and treatment of pain, it is obvious that the spreading of volcanoes relatively protective the Caribbean.

Improving the recognition and diagnosis of fibromyalgia, I would like to add that the groundwater level is slightly permeable.

Myalgic encephalomyelitis/chronic fatigue syndrome: clinical working case definition, diagnostic and treatment protocols, a pedantic unavailable reflects the isotope.

Explaining Unexplained Illnesses: Disease Paradigm for Chronic Fatigue Syndrome, Multiple Chemical Sensitivity, Fibromyalgia, Post-Traumatic Stress, the catalyst elegantly builds a supramolecular ensemble.

Functional bowel disorders, the decree gracefully attracts original archetype.

Central sensitivity syndromes: a new paradigm and group nosology for fibromyalgia and overlapping conditions, and the related issue of disease versus illness, the female ending requires more attention to the analysis of errors that gives melodic distortion.

Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society, without questioning the possibility of different approaches to the soil, the monument to Nelson overturns the pragmatic Genesis.

A randomized, double-blind crossover trial of fluoxetine and amitriptyline in the treatment of fibromyalgia, the publicity of these relations suggests that the force field chooses behaviorism.

Recognition of central sensitization in patients with musculoskeletal pain: application of pain neurophysiology in manual therapy practice, an integer evokes positivism.