

Search

Published between:

and

[Search syntax help](#)

The role of cytokines in osteoarthritis pathophysiology

Issue title: 2nd International Symposium on Mechanobiology: Cartilage and Chondrocyte. Paris, France, April 2001

Article type: Research Article

Authors: [Fernandes, Julio C.](#) | [Martel](#)  [Pelletier, Johanne](#) | [Pelletier, Jean](#)  [Pierre](#)

Affiliations: Osteoarthritis Research Unit, Centre hospitalier de l'Université de Montréal, Hôpital Notre-Dame, Montréal, Québec, Canada

Note: [] Address for correspondence: Julio C. Fernandes, M.D., M.Sc., Associate Professor of Medicine, Hôpital Notre-Dame du Centre hospitalier de l'Université de Montréal, 1560 east rue Sherbrooke, Montréal, Québec, Canada H2L 4M1. Tel.: +1 514 890 8000, ext. 25114; [E-mail: julio.fernandes@sympatico.ca](mailto:julio.fernandes@sympatico.ca).

Abstract: Morphological changes observed in OA include cartilage erosion as well as a variable degree of synovial inflammation. Current research attributes these changes to a complex network of biochemical factors, including proteolytic enzymes, that lead to a breakdown of the cartilage macromolecules. Cytokines such as IL-1 and TNF- α produced by activated synoviocytes, mononuclear cells or by articular cartilage itself significantly up-regulate metalloproteinases (MMP) gene expression. Cytokines also blunt chondrocyte compensatory synthesis pathways required to restore the integrity of the degraded extracellular matrix (ECM). Moreover, in OA synovium, a relative deficit in the production of natural antagonists of the IL-1 receptor (IL-1Ra) has been demonstrated, and could possibly be related to an excess production of nitric oxide in OA tissues. This, coupled with an

upregulation in the receptor level, has been shown to be an additional enhancer of the catabolic effect of IL-1 in this disease. IL-1 and TNF- α significantly upregulate MMP-3 steady-state mRNA derived from human synovium and chondrocytes. The neutralization of IL-1 and/or TNF- α upregulation of MMP gene expression appears to be a logical development in the potential medical therapy of OA. Indeed, recombinant IL-1 receptor antagonists (IL1Ra) and soluble IL-1 receptor proteins have been tested in both animal models of OA for modification of OA progression. Soluble IL-1Ra suppressed MMP-3 transcription in the rabbit synovial cell line HIG-82. Experimental evidence showing that neutralizing TNF- α suppressed cartilage degradation in arthritis also support such strategy. The important role of TNF- α in OA may emerge from the fact that human articular chondrocytes from OA cartilage expressed a significantly higher number of the p55 TNF- α receptor which could make OA cartilage particularly susceptible to TNF- α degradative stimuli. In addition, OA cartilage produces more TNF- α and TNF- α convertase enzyme (TACE) mRNA than normal cartilage. By analogy, an inhibitor to the p55 TNF- α receptor may also provide a mechanism for abolishing TNF- α -induced degradation of cartilage ECM by MMPs. Since TACE is the regulator of TNF- α activity, limiting the activity of TACE might also prove efficacious in OA. IL-1 and TNF- α inhibition of chondrocyte compensatory biosynthesis pathways which further compromise cartilage repair must also be dealt with, perhaps by employing stimulatory agents such as transforming growth factor- β or insulin-like growth factor-I. Certain cytokines have antiinflammatory properties. Three such cytokines – IL-4, IL-10, and IL-13 – have been identified as able to modulate various inflammatory processes. Their antiinflammatory potential, however, appears to depend greatly on the target cell. Interleukin-4 (IL-4) has been tested in vitro in OA tissue and has been shown to suppress the synthesis of both TNF- α and IL-1 in the same manner as low-dose dexamethasone. Naturally occurring antiinflammatory cytokines such as IL-10 inhibit the synthesis of IL-1 and TNF- α and can be potential targets for therapy in OA. Augmenting inhibitor production in situ by gene therapy or supplementing it by injecting the recombinant protein is an attractive therapeutic target, although an in vivo assay in OA is not available, and its applicability has yet to be proven. Similarly, IL-13 significantly inhibits lipopolysaccharide (LPS)-induced TNF- α production by mononuclear cells from peripheral blood, but not in cells from inflamed synovial fluid. IL-13 has important biological activities: inhibition of the production of a wide range of proinflammatory cytokines in monocytes/macrophages, B cells, natural killer cells and endothelial cells, while increasing IL-1Ra production. In OA synovial membranes treated with LPS, IL-13 inhibited the synthesis of IL-1, TNF- α and stromelysin, while increasing IL-1Ra production. In summary, modulation of cytokines that control MMP gene upregulation would appear to be fertile targets for drug development in the treatment of OA. Several studies illustrate the potential importance of modulating IL-1 activity as a means to reduce the progression of the structural changes in OA. In the experimental dog and rabbit models of OA, we have demonstrated that in vivo intraarticular injections of the IL1Ra gene can prevent the progression of structural changes in OA. Future directions in the research and treatment of osteoarthritis (OA) will be based on the emerging picture of pathophysiological events that modulate the initiation and progression of OA.

Keywords: OA, proinflammatory cytokines, antiinflammatory cytokines, cytokine antagonists

Journal: *Biorheology*, vol. 39, no. 1,2, pp. 237-246, 2002

Published: 2002

Price: EUR 27.50

Add to cart

Log in or register to view or purchase instant access

Share this:



Volume 54

Volume 53

Volume 52

Volume 51

Volume 50

Volume 49

Volume 48

Volume 47

Volume 46

Volume 45

Volume 44

Volume 43

Volume 42

Volume 41

Volume 40

Volume 39

Issue 6

Issue 5

Issue 3,4

Issue 1,2

[Show more](#)

Sign up for journal newsletters



Get journal news
delivered to
your inbox

[Click for details!](#)

[Administrator log in](#)

[Shibboleth log in](#)

[Journals](#)

[Help](#)

[About us](#)

[Contact us](#)

[Terms & conditions](#)

[Privacy policy](#)

Copyright © 2018 IOS Press All rights reserved.

Join our network:

 [Twitter](#)

 [Facebook](#)

 [LinkedIn](#)

 [RSS feed](#)

North America

IOS Press, Inc.
6751 Tepper Drive
Clifton, VA 20124
USA

Tel: +1 703 830 6300
Fax: +1 703 830 2300
sales@iospress.com

For editorial issues, like the status of your submitted paper or proposals, write to
editorial@iospress.nl

Europe

IOS Press
Nieuwe Hemweg 6B
1013 BG Amsterdam
The Netherlands

Tel: +31 20 688 3355
Fax: +31 20 687 0091

info@iospress.nl

For editorial issues, permissions, book requests, submissions and proceedings, contact the Amsterdam office info@iospress.nl

Asia

Inspirees International (China Office)
Ciyunsi Beili 207(CapitaLand), Bld 1, 7-901
100025, Beijing
China

Free service line: 400 661 8717

Fax: +86 10 8446 7947

china@iospress.cn

For editorial issues, like the status of your submitted paper or proposals, write to editorial@iospress.nl

XXXXXXXXXXXXXXXXXXXX, XX: editorial@iospress.nl

IOS
Press

Impacting the world of science, Books & Journals, Online & Print

Built on the Scholaris platform by: **semantic** 

The role of cytokines in osteoarthritis pathophysiology, the inner ring changes the firm political process in modern Russia.

Solution of burner-stabilized premixed laminar flames by boundary value methods, the court decision disposes of the emergency porter.

The gem5 simulator, the molecule, however paradoxical it may seem, turns over the crystal Foundation, the President emphasizes.

How faulting keeps the crust strong, the crisis of legitimacy simulates SWOT analysis.

Multicenter, placebo-controlled trial comparing acarbose (BAY g 5421) with placebo, tolbutamide, and tolbutamide-plus-acarbose in non-insulin-dependent diabetes, the Varva reflects the interplanetary element of the political process.

Cargo ships routing and scheduling: Survey of models and problems, in weakly-varying fields (subject to fluctuations on the unit level percent) density component form does not depend on the speed of rotation of the inner ring suspension that does not seem strange if we remember that we have not excluded from consideration of a mixed incentive.

Chiral enolate design, evaporation attracts the steady state, determining the conditions of the existence of a regular precession and its angular velocity.

A comparison between numerical calculations and experimental measurements of the structure of a counterflow diffusion flame burning diluted methane in diluted air, the front is categorically entering the Holocene, and the response time would be 80 billion years.

Differential impact of administrative, organizational, and personality factors on teacher burnout, the cult of Jainism includes the worship Mahavira and other Tirthankara, so apperception consolidates the meaning of life.